



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Statistical reports of States and cities of the United States—Yearly and monthly.

CALIFORNIA—*San Diego*.—Month of July, 1902. Estimated population, 20,000. Total number of deaths, 25, including 8 from tuberculosis.

MARYLAND—*Baltimore*.—Month of July, 1902. Estimated population, 525,000—white, 445,000; colored, 80,000. Total number of deaths, 1,052—white, 817; colored, 235—including enteric fever, 19; measles, 8; scarlet fever, 5; whooping cough, 13; diphtheria, 2, and 87 from tuberculosis.

MASSACHUSETTS—*Fitchburg*.—Month of July, 1902. Estimated population, 33,848. Total number of deaths, 22, including 1 from measles.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended August 23, 1902, from 68 observers, indicate that cholera infantum, dysentery, enteric fever, erysipelas, remittent fever, pneumonia, and diphtheria were more prevalent, and measles was less prevalent than in the preceding week. Meningitis was reported present at 7, diphtheria at 16, whooping cough at 18, measles at 23, smallpox at 36, scarlet fever at 48, enteric fever at 86, and phthisis pulmonalis at 219 places.

MISSOURI—*St. Louis*.—Month of July, 1902. Census population, 621,000—white, 582,000; colored, 39,000. Total number of deaths, 910—white, 796; colored, 114—including diphtheria, 15; measles, 3; scarlet fever, 9; whooping cough, 25, and 135 from tuberculosis.

NEW JERSEY—*Hudson County*.—Month of July, 1902. Estimated population, 409,810. Total number of deaths, 794, including enteric fever, 9; diphtheria, 13; measles, 3; scarlet fever, 5; whooping cough, 11; smallpox, 16, and 71 from tuberculosis.

Passaic.—Four weeks ended August 23, 1902. Census population, 27,777. Total number of deaths, 62, including diphtheria, 1, and 1 from enteric fever.